

# Rift Valley Fever Symptoms

**Texas A&M University  
College of Veterinary Medicine**

**University of Pretoria  
Department of  
Veterinary Tropical Diseases**

Jeffrey Musser, DVM, PhD  
Suzanne Burnham, DVM

Professor J.A.W. Coetzer



# Special thanks for materials borrowed with permission from presentations by:

- Dr Linda Logan, “Rift Valley Fever” CSU Foreign Animal Disease Training Course, College of Veterinary Medicine and Biomedical Sciences, August 1-5, 2005.
- Professor JAW Coetzer, Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, “Rift Valley Fever” presented at the FEAD course in Knoxville, Tenn. 2005.





# Rift Valley Fever

*In this presentation the authors especially drew from the first hand experience of their colleagues in South Africa. Personal interviews as well as standard research sources provide the insights we bring you for the recognition of this exotic disease.*

**JAW Coetzer**  
**Jeffrey Musser**  
**Suzanne Burnham**



Rift Valley Fever- 2006



# Rift Valley Fever

- Rift Valley Fever (RVF) is an arthropod-borne, acute, fever-causing viral disease of sheep, goats, cattle and people.





Rift Valley fever in Africa causes abortions in sheep, cattle and goats high mortalities in lambs and kids and generalized disease in man.



Rift Valley Fever- 2006



# Rift Valley Fever

RVF was first observed when European stocks of domestic animals were introduced to Africa. These species are more severely affected than native African stock.



Rift Valley Fever- 2006



# Host Range

- Mainly a disease of sheep



# Host Range

- In Sheep  
Mortality in lambs  
under 2 weeks of  
age approaches  
100%

Mortality in older sheep  
reaches 30%  
with abortions  
approaching 100%





# Host range

- Cattle are less susceptible than sheep, some are subclinical; mortality averages 5% with some abortions



# Host Range

- Goats
- Buffalo



Rift Valley Fever- 2006



# Host Range

- Domestic dogs and cats – susceptible but usually only have asymptomatic viremia
- Swine - resistant
- Birds - refractory, no virus isolation



# Host Range

- Horses – have viremia but are resistant



# Host range - wildlife

- Springbok
- African Buffalo
- Camels (in Egypt)
- Water buffalo in Egypt



# Host range - wildlife



Water buffalo - up to 50% abortion rate



Rift Valley Fever- 2006



# Host range - wildlife



Camels (in Egypt) - inapparent disease except abortions



Rift Valley Fever- 2006



## Rift Valley fever host range and disease severity


Mortality ~ 100%	Severe Illness Abortion, Low Mortality	Severe Illness Viremia Abortion	Infection Viremia	Refractive to infection
Lambs Calves Kids Puppies Kittens White mice Hamster Field mice Door mice Field voles	Sheep Cattle Goats Water buffalo  Humans	Monkeys Camels Rats Gray squirrels	Horses Cats Dogs Monkeys	Guinea pigs Rabbits Pigs Hedgehogs Tortoises Frogs Chickens Canaries Pigeons Parakeets





# Incubation period

- 1-6 days
- 12-36 hours in lambs; will be dead before they can acquire passive immunity



# Clinical signs

## Sheep and Goats

- Incubation period less than 3 days
- High rate of abortion at any stage of gestation
- Some show no symptoms





In pregnant ewes, abortion may approach 100%  
Aborted fetus is usually autolyzed.



Rift Valley Fever- 2006



# Clinical signs Sheep and Goats

- Abortion rate in sheep from 40 – 100%
- Ewe may also retain the placenta
- Endometritis is another complication after aborting the fetus



USDA



Rift Valley Fever- 2006



# Clinical signs

## Sheep and Goats

### Early signs

- Fever 40-41°C
- Loss of appetite
- Jaundice
- Weakness



# Clinical signs Sheep and Goats

- Encrustation around the muzzle from bloody nasal discharge



# Clinical signs Sheep and Goats



- Some develop diarrhea



Rift Valley Fever- 2006



# Clinical signs Sheep and Goats

- Acute death may occur  
in 20-30% of adults





# Clinical signs Sheep and Goats



Heavy sheep losses  
occur  
during epidemic



Rift Valley Fever- 2006



# Clinical Signs in lambs and kids

■ Newborn Lambs, Kids: Most severe in young lambs under 2wks old (mortality has high as 90%)

- fever (40-42°C),
- anorexia,
- weakness,



# Clinical Signs in lambs and kids



Lambs seem reluctant to move;  
they have signs of abdominal pain,  
rapid respiration and listlessness.



# Clinical Signs in lambs and kids

Death may occur within 24 to 36 hours after the first signs appear. Death is due to severe liver necrosis and vascular collapse.



# Clinical signs in cattle



- Anorexia

- Weakness

- Fetid diarrhea

- Often only sign is a drop in calving rates



# Clinical signs in cattle

- Calves: fever (40-41°C), depression. Mortality rate: 10-70% Death occurs about 2-8 days after the first signs appear.
- Adults: fever (40-41°C), excessive salivation, anorexia, weakness, fetid diarrhea, fall in milk yield. Abortion may reach 85% in the herd. Mortality rate is usually less than 10%



# Clinical signs in cattle



Rift Valley Fever- 2006



# Clinical signs in cattle

- Disease most severe signs are seen in young animals
- Symptoms may be prolonged and will include jaundice in some calves
- Aborted calves are moderately autolyzed.





# Clinical signs

- Dogs: Abortions may occur in adult dogs; severe disease and death usually only in puppies
- Cats: Death in kittens



# Relative susceptibility

Newborn ruminants	++++
Pregnant ruminants	++++
Sheep and young cattle	+++
Adult cattle, goats, sheep	++
Humans	++
Dogs, cats and camels	+
Pigs	-



# Transmission

- RVF is primarily transmitted from animal to animal by a mosquito



- *Aedes*, *Culex*, *Anopheles*, *Erethmapodites*, *Monsosmia*



# Necropsy findings

- Massive hepatitis: hemorrhages, necrotic foci, marked enlargement, orange-brown, friable, edematous liver tissue

*(“If you open a newborn lamb, the liver jumps into your face” Coetzer)*

- Chocolate-brown digested blood in abomasum, hemorrhages in intestinal mucosa, free blood in lumen



# Pathology Summary

- Focal or generalized hepatic necrosis
- Congestion, enlargement, and discoloration of liver with subcapsular hemorrhages
- Brown-yellowish color of liver in aborted fetuses
- Hemorrhagic enteritis
- Icterus (low percentage)
- Widespread cutaneous hemorrhages, petechial to ecchymotic hemorrhages on parietal and visceral serosal membranes
- Enlargement, edema, hemorrhages and necrosis of lymph nodes
- Congestion and cortical hemorrhages of kidneys and gallbladder



# Necropsy : new-born lambs



Liver massively enlarged; hemorrhages; orange-brown color; small areas of necrosis. The liver is very friable.



# Necropsy : new-born lambs



Gall bladder hemorrhage; Abomasum diffuse hemorrhage,  
serosa has petechial hemorrhage



# Necropsy : new-born lambs





Abomasum shows diffuse chocolate brown hemorrhages, serosa has petechial hemorrhages, necrotic foci, and D. I. C.





# Necropsy : adult sheep



May look like plant poisoning



Rift Valley Fever- 2006



# Necropsy : adult sheep




Gall bladder contains frank hemorrhage



Rift Valley Fever- 2006



# Necropsy : sheep & cattle




Abomasum is edematous similar to Heartwater



Rift Valley Fever- 2006



# Necropsy : adult cattle



Close up of gall bladders



# Necropsy : adult cattle



Petechia



Rift Valley Fever- 2006



# Necropsy : cattle



Spleen with many hemorrhages



Rift Valley Fever- 2006



# Necropsy : other lesions

Enlarged  
lymph nodes



Rift Valley Fever- 2006



# Differential Diagnosis

- Abortifacient agents
- Agents causing hepatitis
- Agents that cause hemorrhages





# Differential Diagnosis

- Bluetongue
- Wesselsbron disease
- Enterotoxemia of sheep
- Ephemeral fever
- Brucellosis
- Vibriosis
- Trichomonosis



# Differential Diagnosis

- Nairobi sheep disease
- Heartwater
- Ovine enzootic abortion
- Toxic plants
- Bacterial septicemias (Pasteurella, Salmonella, Anthrax)
- Rinderpest and Peste des petits ruminants



# Suspect Rift Valley Fever if:



- High mortalities in lambs, kids and calves following increase in mosquito populations
- Disease is milder in adults than in newborns
- Abortions in sheep, goats and cattle
- Extensive necrotic liver changes
- Influenza symptoms in people working with sick animals or handling infected carcasses



# Rift Valley Fever - Bibliography

1. Linda L Logan, DVM PhD, USDA APHIS Attaché, North Africa, East Africa, Middle East, “Rift Valley Fever” CSU Foreign Animal Disease Training Course, Aug 1-5, 2005.
2. Professor J A W Coetzer, Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, “Rift Valley Fever”
3. USAHA, Foreign Animal Diseases, 1992 Edition, p.311-317
4. W.A. Geering, A.J. Foreman and M.J. Nunn, Exotic Diseases of Animals, 1995 Australian Govt Publishing Service, Canberra; p.218- 224.





An excellent video about Rift Valley Fever is available from:  
[http://www.up.ac.za/academic/veterinary/depts\\_vtd\\_teach/index.htm](http://www.up.ac.za/academic/veterinary/depts_vtd_teach/index.htm)



# Image Watermark

- “KAW” images by Dr. Ken A. Waldrup
- “Coetzer” images used with permission by Dr. Koos Coetzer
- “LLogan” images by Dr. Linda Logan
- “Suz” images by Dr Suzanne Burnham
- “MFitilodze” images by Dr. M. W. (Bill) Mfitilodze Bunda College of Agriculture, University of Malawi



# Acknowledgements

Special thanks to

Professor JAW Coetzer

Linda Logan, DVM PhD, USDA

Ken Waldrup, DVM, PhD

Robin Sewell, DVM

Kelsey Pohler- Research Assistant

